



McCabe, R., Sterno, I., Priebe, S., Barnes, R., & Byng, R. (2017). How do healthcare professionals interview patients to assess suicide risk? *BMC Psychiatry*, 17, [122]. <https://doi.org/10.1186/s12888-017-1212-7>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1186/s12888-017-1212-7](https://doi.org/10.1186/s12888-017-1212-7)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via BioMed Central at <https://bmcpsy psychiatry.biomedcentral.com/articles/10.1186/s12888-017-1212-7>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Transcription conventions

.hhh	Audible inhalation
hhh	Audible exhalation
:	Extended sound
↑	Rising intonation
↓	Falling intonation
?	Rising inflection
_____	Emphasis (word or part of word underlined)
◦ ◦	Talk is quieter than the surrounding talk
< >	Talk is faster than the surrounding talk
UPPERCASE	Talk is louder than the surrounding talk
!	Animated tone
=	Latched utterance, no interval between utterances
[]	Beginning and end of overlapping talk
()	Transcriptionist doubt
(.)	A pause of less than 0.2 seconds
(0.0)	Silence measured in seconds and tenths of seconds